



## Browning Seed, Inc.

3101 Interstate 27 Frontage  
Plainview, TX 79072

Phone: (806) 293-5271

Website: [www.browningseed.com](http://www.browningseed.com)

### Positioning/Management

Cheyenne can be used for both grazing and hay production throughout the spring and summer months under good management practices. Razed, apply up to 150 lbs. of nitrogen per year in split applications throughout the summer. Last fertilizer application each year should be done 4 to 6 weeks before a killing frost to increase cold tolerance. Delay grazing until forage is 8" to 10" tall. Do not graze or clip for hay shorter than 2". Rotate animals more often during periods of drought stress.

**This is a certified PVP variety of  
Seeds West, Inc.**

## Cheyenne II

### Forage Bermudagrass Seed

### Adaptation

Across the southern 1/3 of the U.S. from California to Virginia. Particularly well suited to sandy soils but will thrive in most well-drained loam soils.

### Planting

- Rate: 15 lbs./acre
- Date: Late spring through early summer when soil temperatures are 65 degrees fahrenheit or above. Plow and cultipack to develop a firm seedbed. Proper firmness is indicated by a heel print no more than 1/8" deep in the soil.
- Depth: 1/8" deep maximum.
- Fertilizer: Due to varying soil conditions and types it is necessary to conduct a soil test each year in order to supply the proper nutrients. Apply 20-30 lbs. of nitrogen at planting time. When the new plants start to run, apply 50-60 lbs. of nitrogen. After the stand is established, apply 70-100 lbs. of nitrogen after each cutting.

***Browning Seed - "Programmed for Profit"***