



# Browning Seed, Inc.

3101 Interstate 27 Frontage  
Plainview, TX 79072

Phone: (806) 293-5271

Website: www.browningseed.com

## Agronomic Attributes

Potential Forage Yield.....	4
Early Growth Rate.....	3
Regrowth.....	3
Average Plant Height.....	6'-7'
Standability.....	3
Drought Tolerance.....	3
Leafiness.....	3
Avg. Seeds/Lb. ....	18,000
Fast Dry Down.....	1
Sweetness.....	2

## Primary Uses

Hay.....	1
Hay Plow Down.....	2
Silage.....	3
Grazing/Pasture.....	1
Greenchop.....	2

## Feed Analysis

Crude Protein.....	8.94
Acid Detergent Fiber.....	29.48
Total Digestible Nutrients.....	67.05
NE (lactation) Mcal/lb.....	0.69
NE (gain) Mcal/lb.....	0.67

## Yield

1 to 3 cuttings dependent upon planting date, availability of water, and growing season. 3 to 6 tons/acre dry matter (20 to 40 tons green weight).

**- HIGHLY NUTRITIOUS-**

## Planting Rate (LBS./ACRE)

Optimal Planting Method		
	Wide Rows	Narrow Rows
Moisture	30" to 40"	Drilled or Broadcast
Adequate	15 to 30 LBS/AC	50 to 80 LBS/AC
Limited	12 to 20 LBS/AC	30 to 60 LBS/AC

# Bundle King

## Sterile Forage Sorghum

## Positioning/Management

Bundle King is a fast growing medium early hybrid that produces very sweet, juicy stalks that can be either grazed, baled or used for greenchop. Bundle King is a sterile hybrid that will not produce seed in the absence of pollen, which allow for more of the proteins and sugars to stay in the plant instead of being transferred to the head in the growth process. Bundle King also has an excellent leaf to stem ratio and is great to start new calves on.

## Annual Summer Management

**Planting Date:** Late April through mid August. Soil temperature should be 60-65 or warmer. Planting too early could cause slow early growth and reduction in population.

**Planting Depth:** 3/4" to 1 1/2" deep dependent upon the soil moisture.

**Row Width:** Preferably drilled in narrow rows .

**Planting Method:** Drill preferred.

**Fertility:** Nitrogen: 75 to 100 lbs./acre PPI & 50 to 75 lbs./acre after first cutting. Phosphorus: 20-60 lbs./acre, more if the soil testing indicates less than 30 lbs./acre. Potassium: 40-80 lbs./acre. A soil test should be performed for better accuracy. The above fertilization rates serve as a guide only.

## Suggested Harvest Management

Do not cut or graze until the forage reaches 18" in height. Cut at 36"- 48" or 60 -75 days for the best possible quality and highest possible yields. Cut 6"- 8" or above the second internode with a crimper to speed the drying and regrowth process. Harvest forage at about 70 -75% moisture. Chop at 1/4" -1/8" to get optimal packing.

## Footnotes: 1- Numerical Rating: 1 to 9

- 1= Excellent
- 5= Average
- 9= Poor

Ratings and descriptions are based on the research and field observations that were compared with products from multiple locations and years. Read all labels and bag tags due to the reason that they contain conditions of sale, including limitations of warranty and remedy.

**- "Super Sweet" -**

**Browning Seed - "Programmed for Profit"™**