



Browning Seed, Inc.

3101 Interstate 27 Frontage

Plainview, TX 79072

Phone: (806)293-5271

Website: www.browningseed.com

Tridan®

A Three Way Cross

Sorghum X Sudangrass Hybrid

Agronomic Attributes

Potential Forage Yield.....	2
Early Growth Rate.....	2
Regrowth.....	1
Average Plant Height.....	5'-8"
Standability.....	3
Drought Tolerance.....	2
Leafiness.....	3
Avg. Seeds/Lb.	18,000
Fast Dry Down.....	1
Protein Content.....	High

Primary Uses

Hay.....	1
Hay Plow Down.....	1
Silage.....	1
Grazing/Pasture.....	2
Greenchop.....	4

Suggested Harvest Height

Hay/Plow Down.....	Pre-boot or Flag Stage or 40"-60"
Grazing.....	30-45 days or 20"-25"
Silage.....	35"-50"
Greenchop.....	24"-40"

Yield

2 to 3 cuttings dependent upon planting date, availability of water, and growing season. 3 to 7 tons/acre dry matter (15 to 20 tons green weight).

"Sugar Sweet"®



"The Livestock Dessert"

Positioning/Management

Tridan® is a sweet, leafy, highly nutritious forage sorghum hybrid. This full season multipurpose sorghum sudangrass hybrid produces excellent grazing, hay, green chop, silage or green ever crop. Tridan® will furnish summer grazing of up to five cows per acre with good management practices; and it recovers quickly after grazing or cutting. Cattle thrive on lush forage during the summer and sweet hay in the winter.

Annual Summer Management

Planting Date: Late April through mid August. Soil temperature should be 60-65 or warmer. Planting too early could cause slow early growth and reduction in population.

Planting Depth: 3/4" to 1 1/2" deep dependent upon the soil moisture.

Row Width: Preferably drilled in narrow rows.

Planting Method: Drill preferred.

Fertility: Nitrogen: 75 to 100 lbs./acre PPI & 50 to 75 lbs./acre after first cutting. Phosphorus: 20-60 lbs./acre, more if the soil testing indicates less than 30 lbs./acre. Potassium: 40-80 lbs./acre. A soil test should be performed for better accuracy. The above fertilization rates serve as a guide only.

Suggested Harvest Management

Do not cut or graze until the forage reaches 18" in height. Cut at 36"- 48" or 60 -75 days for the best possible quality and highest possible yields. Cut 6"- 8" or above the second internode with a crimper to speed the drying and regrowth process. Harvest forage at about 70 -75% moisture. Chop at 1/4" -1/8" to get optimal packing.

Footnotes: 1- Numerical Rating: 1 to 9

- 1= Excellent
- 5= Average
- 9= Poor

Ratings and descriptions are based on the research and field observations that were compared with products from multiple locations and years. Read all labels and bag tags due to the reason that they contain conditions of sale, including limitations of warranty and remedy.

Browning Seed - "Programmed for Profit"