



Browning Seed, Inc.

3101 S. IH 27
Plainview, TX 79072
Phone: (806) 293 – 5271
Website: www.browningseed.com



Agronomic Attributes

Potential Forage Yield	2
Early Growth Rate	3
Regrowth	3
Average Plant Height	6' - 10'
Standability	1
Drought Tolerance	2
Leafiness	3
Avg. Seeds/ Lb.	18,000 – 20,000
Fast Dry Down	2
Lignin Content	Low

Primary Uses

Hay	2
Hay Plow Down	1
Silage	1
Grazing/Pasture	3
Greenchop	3

Suggested Harvest Height

Hay/Plow Down	Pre-boot, Flag Leaf Stage, or 8' – 10'
Grazing	30-45 Days or 18" - 24"
Silage	97"- 120"
Greenchop	24" – 120"

Yield

1 - 2 Cuttings dependent upon planting date, availability of water, and growing season. 3 – 6 tons/acre dry matter (15 to 20 tons green weight).

Planting Rate (LBS./Acre)

	Optimal Planting Method	
	Wide Rows	Narrow Rows
Moisture Level	30" to 40"	Drilled or Broadcast
Adequate	20 to 50 LBS/AC	50 to 70 LBS/AC
Limited	10 to 30 LBS/AC	30 to 50 LBS/AC

Headless Wonder™ PPS BMR is a Photoperiod Sensitive Sorghum/Sudangrass Hybrid.

Positioning/ Management

Headless Wonder™ PPS BMR has a good regrowth capacity, ideal for intensive grazing. Has good genetic potential, yielding 8,000 to 12,000 lbs per acre of high-quality forage. For its photoperiod sensitive trait, it delays heading out.

Annual Summer Management

Planting Date: Late April through mid August. Soil temperature should be 65-70 or warmer. Planting too early could cause slow early growth and reduction in population.

Planting Depth: 3/4" to 1 1/2" deep dependent upon the soil moisture.

Row Width: Preferably drilled in narrow rows.

Planting Method: Drill preferred.

Fertility: Nitrogen: 75 to 100 lbs./acre PPI & 50 to 75 lbs./acre after first cutting. Phosphorus: 20-60 lbs./acre, more if the soil testing indicates less than 30 lbs./acre. Potassium: 40-80 lbs./acre. A soil test should be performed for better accuracy. The above fertilization rates serve as a guide only. We highly recommend soil testing and consultation of a soil scientist for best results.

Suggested Harvest Management

Do not cut or graze until the forage reaches 18" in height. Cut at 36"- 48" or 60 -75 days for the best possible quality and highest possible yields. Cut 6"- 8" or above the second internode with a crimper to speed the drying and regrowth process. Harvest forage at about 70 -75% moisture. Chop at 1/4" -1/8" to get optimal packing.

Footnotes: 1– Numerical Rating: 1 to 9

1= Excellent

5= Average

9= Poor

Ratings and descriptions are based on the research and field observations that were compared with products from multiple locations and years. Read all labels and bag tags due to the reason that they contain conditions of sale, including limitations of warranty and remedy.