



BROWNING SEED, INC.

3101 S. IH 27 PLAINVIEW, TX79072

PHONE: (806) 293-5271

WEBSITE: WWW.BROWNINGSEED.COM



Agronomic Attributes

Potential Forage Yield	4
Early Growth Rate	3
Regrowth	3
Average Plant Height	6' - 7'
Standability	3
Drought Tolerance	3
Leafiness	3
Avg. Seeds/ Lb.	18,000
Fast Dry Down	1
Sweetness	2

Primary Uses

Hay	1
Hay Plow Down	2
Silage	3
Grazing/Pasture	1
Greenchop	2

Feed Analysis

Crude Protein	8.94
Acid Detergent Fiber	29.48
Total Digestible Nutrients	67.05
NE (Lactation) Mcal/lb	0.69
NE (Gain) Mcal/lb	0.67

Planting Rate (LBS/ Acre)

	Optimal Planting Method	
	Wide Rows	Narrow Rows
Moisture Level	30" to 40"	Drilled or Broadcast
Adequate	15 to 30 LBS/AC	50 to 80 LBS/AC
Limited	12 to 20 LBS/AC	30 to 60 LBS/AC

Footnotes: 1- Numerical Rating: 1 to 9 1= Excellent 5= Average 9= Poor

Ratings and descriptions are based on the research and field observations that were compared with products from multiple locations and years. Read all labels and bag tags due to the reason that they contain conditions of sale, including limitations of warranty and remedy.

Why our Bundle King?

Bundle King is a fast-growing, medium-early hybrid that stands out as an excellent choice for customers. Its appeal lies in the production of exceptionally sweet and juicy stalks, which can be utilized in various ways such as grazing, baling, or greenchop. The fact that Bundle King is a sterile hybrid is a significant advantage as it prevents seed production in the absence of pollen. This unique trait allows more proteins and sugars to remain in the plant, enhancing its nutritional value. Furthermore, Bundle King showcases an impressive leaf to stem ratio, ensuring a well-balanced and nutrient-rich feed. This makes it an ideal option for starting new calves and catering to their specific dietary requirements.

Annual Summer Management

Planting Date: Late April through mid August. Soil temperature should be 60-65 or warmer. Planting too early could cause slow early growth and reduction in population.

Planting Depth: 3/4" to 1 1/2" deep dependent upon the soil moisture.

Row Width: Preferably drilled in narrow rows.

Planting Method: Drill preferred.

Fertility: Nitrogen: 75 to 100 lbs./acre PPI & 50 to 75 lbs./acre after first cutting. Phosphorus: 20-60 lbs./acre, more if the soil testing indicates less than 30 lbs./acre. Potassium: 40-80 lbs./acre. A soil test should be performed for better accuracy.

The above fertilization rates serve as a guide only.

Suggested Harvest Management

Do not cut or graze until the forage reaches 18" in height. Cut at 36"-48" or 60 -75 days for the best possible quality and highest possible yields. Cut 6"- 8" or above the second internode with a crimper to speed the drying and regrowth process. Harvest forage at about 70 - 75% moisture. Chop at 1/4" -1/8" to get optimal packing.

Yield

1 to 3 cuttings dependent upon planting date, availability of water, and growing season. 3 to 6 tons/acre dry matter (20 to 40 tons green weight).

